

BLEPHAROPLASTY/EYELID SURGERY

Post-operative Instructions Following Blepharoplasty

Ramin A. Behmand, MD
1776 Ygnacio Valley Road, Suite 108
Walnut Creek, CA 94598
Tel: (925)-939-9200

Call the office immediately if you have:

- A temperature of 101 or greater
- Redness that is beginning to spread from the incision site
 - Any unusual or painful swelling and redness
 - Any active bleeding soaking more than a 4x4 gauze
- Any purulent “pus like” drainage coming from the incision site
 - Pain that is not relieved by your pain medications

1. Right after Surgery: Your main goal during the first few days to a week after surgery is to help minimize swelling. You can do this by keeping your head elevated at a 45 degree angle, either by sleeping in a recliner or having several pillows behind your back. Your eyes might be uncomfortable. If you have excessive pain or loss of vision you should call the office at (925)939-9200 immediately.

2. Swelling and bruising: Swelling will be at its most noticeable stage at 2-3 days after surgery and will then start to subside. Most bruising disappears by around 7-14 days after surgery.

3. Redness: Expect the white of your eye to be red. This may even get worse over the next week but will clear. Do not rub or bump your eyes.

4. Eating: Start with light & soft meals. Avoid chewy/fatty foods during the first 24 hours after surgery. As you get more comfortable, you may progress to regular food. Drinking fluids is very important after surgery; at least 1 quarts a day (about 1 liters). Try to drink non-caffeinated fluids.

5. Ice: Use an ice mask as much as possible in the first 24 hours only and make sure that you use a clean cloth as a barrier between your skin and the ice pack.

6. Medications: All your medications will be prescribed for you during your preoperative visit at our office. Please fill these prescriptions prior to your surgery. Take your medications as prescribed. Try not to take pain medication on an empty stomach; it can cause nausea. You only need to take the pain medication, if you have discomfort. Taper off the pain medication when your pain level starts to decrease during the first week. The stool softener helps prevent and relieve constipation, but it is effective only if you are drinking plenty of fluids with it.

7. Driving: Do not drive yet. Once you are NO LONGER TAKING pain medication at all, and are feeling comfortable, you may start driving.

8. Incision Care: Use Bacitracin or Polysporin (antibiotic ointment) 3 times daily on the incisions to minimize scabbing.

9. Sun Exposure: Avoid direct sun exposure for at least 6 weeks after surgery. Wear sunscreen at all times during the day (once you are given the approval to use skin products from Dr. Behmand, usually around postoperative day 10).

10. Exercise: No exercise for 4 weeks after surgery. Avoid bending over, heavy lifting or any other activity that makes your face feel pressure.

11. Skin Care/Makeup: Do not use your regular skincare products OR makeup until your first postoperative appointment. At that time, Dr. Behmand will assess to see if you are ready to start using products again.

12. Contact Lenses: If you wear contact lenses, you can generally start wearing them 3-4 days after surgery depending on your level of comfort. If you have any questions about this, please contact the office.

13. Showering: You may shower 3 days after surgery or when you feel ready. However, do not get water in your eyes for one week after surgery.

14. Follow-up: During your preoperative appointment, you will be given a printout detailing the dates and times for your surgery and your first postoperative follow-up appointment. This sheet will be in the black folder given to you at our office.

15. 24 Hour Answering Service: If anything is urgently needed after 5pm or on the weekend, please be aware that we do have a 24 hour answering service. You may call our regular office line at (925) 939- 9200 to reach the answering service who will be able to transfer you to Dr. Behmand, if need be.